

COUNTEREXAMPLE TO THE TROTTER PRODUCT FORMULA FOR PROJECTIONS

MÁTÉ MATOLCSI AND ROMAN SHVIDKOY

ABSTRACT. We constructed a unitary semigroup $(e^{tA})_{t \geq 0}$ on a Hilbert space and an orthogonal projection P such that the limit

$$\lim_{n \rightarrow \infty} \left[e^{\frac{t}{n}A} P \right]^n$$

does not exist strongly. A similar example with a positive contractive semigroup and positive contractive projection on L_p is also constructed.

1. INTRODUCTION

In this short note we construct counterexamples to the following conjecture.

Suppose P is an orthogonal projection in a Hilbert space H and $(e^{tA})_{t \geq 0}$ is a contractive semigroup on H . Does the limit

$$(1) \quad \lim_{n \rightarrow \infty} \left[e^{\frac{t}{n}A} P \right]^n$$

exist strongly, for all $t > 0$? Or, let $(e^{tA})_{t \geq 0}$ is positive contractive on L_p , for $1 < p < \infty$, and P is a positive contractive projection. Is the same assertion true?

Even though the limit (1) fails to exist in the above settings, there are a few important cases when (1) does make sense and defines a degenerate semigroup. Such is, for example, the case when A is associated with a sectorial sesquilinear form. The reader will find the detailed exposition of the subject and further references in [1], where, in particular, the conjecture was originally posed.

Date: February 1, 2008.

1991 Mathematics Subject Classification. 47d03.

Key words and phrases. Trotter product formula, unitary semigroups, positive contractive semigroups.

The research of the second author was supported by the student part of the NSF grant DMS-9800027.

2. COUNTEREXAMPLES

2.1. Hilbert case. Let us remark that by using the theory of unitary dilations of contractive semigroups in Hilbert spaces one can reduce the first question of the conjecture to the case of unitary semigroups. Therefore, we are looking for a counterexample among unitary semigroups instead of arbitrary contractive ones.

We carry out our construction in the space $L_2[0, 1]$. As an example of unitary semigroup we take the semigroup of multiplication by e^{ith} , where h is a real-valued, measurable function on $[0, 1]$, to be specified later. We choose P to be the one-dimensional orthogonal projection onto the space of constant functions, i.e. $Pf = \mathbf{1} \cdot \int_0^1 f(x)dx$. As a test function on which (1) will fail for $t = 1$, we take $\mathbf{1}$.

Denoting $c_n = \int_0^1 e^{i\frac{1}{n}h(x)}dx$, the function $\left[e^{\frac{1}{n}A}P\right]^n(\mathbf{1})$ becomes $c_n^{n-1}e^{i\frac{1}{n}h}$. However, by the Lebesgue Dominated Convergence Theorem, $\lim_{n \rightarrow \infty} c_n = 1$ as well as $\lim_{n \rightarrow \infty} e^{i\frac{1}{n}h} = \mathbf{1}$ in $L_2[0, 1]$. So, $\lim_{n \rightarrow \infty} \left[e^{\frac{1}{n}A}P\right]^n(\mathbf{1})$ exists in $L_2[0, 1]$ if and only if the numerical limit

$$(2) \quad \lim_{n \rightarrow \infty} c_n^n$$

exists. Now we specify the function h , for which we prove that (2) diverges. Put $h = \sum_{k=1}^{\infty} \chi_{(1/2^k, 1/2^{k-1}]} 2^k \pi$. Then $c_n = \sum_{k=1}^{\infty} \frac{1}{2^k} e^{i\frac{1}{n}2^k \pi}$. We show the following two inequalities

$$(3) \quad \liminf_{n \rightarrow \infty} |c_{2^n}|^{2^n} \geq e^{-(4 + \frac{\pi^2}{4})}$$

$$(4) \quad \limsup_{n \rightarrow \infty} |c_{2^n}|^{2^n} \leq e^{-(6 + \frac{\pi^2}{6} - \frac{\pi^4}{27 \cdot 24 \cdot 7})}.$$

Noticing that $4 + \frac{\pi^2}{4} < 6 + \frac{\pi^2}{6} - \frac{\pi^4}{27 \cdot 24 \cdot 7}$ we get the desired result.

Let us show (3) first. Observe that

$$c_{2^n} = \sum_{k=1}^{n-1} \frac{1}{2^k} e^{i\frac{2^k}{2^n} \pi} - \frac{1}{2^n} + \sum_{k=n+1}^{\infty} \frac{1}{2^k} = \sum_{k=1}^{n-1} \frac{1}{2^k} e^{i\frac{2^k}{2^n} \pi}.$$

Using the inequality $\cos(\alpha) \geq 1 - \frac{\alpha^2}{2}$ we get

$$\begin{aligned} |c_{2^n}| &\geq |\Re c_{2^n}| = \sum_{k=1}^{n-2} \frac{1}{2^k} \cos\left(\frac{2^k}{2^n} \pi\right) \geq \sum_{k=1}^{n-2} \frac{1}{2^k} \left(1 - \frac{\pi^2}{2} \frac{4^k}{4^n}\right) \\ &= 1 - \frac{4}{2^n} - \frac{\pi^2}{2} \frac{1}{4^n} (2^{n-1} - 2) = 1 - \frac{1}{2^n} \left(4 + \frac{\pi^2}{4}\right) + \frac{\pi^2}{4^n}. \end{aligned}$$

Since $\lim_{N \rightarrow \infty} (1 + \frac{a}{N} + \frac{b}{N^2})^N = e^a$, we obtain (3).

To prove (4) let us simplify $c_{2^n 3}$. We have

$$\begin{aligned} c_{2^n 3} &= \sum_{k=1}^{n-1} \frac{1}{2^k} e^{i \frac{2^k}{2^n 3} \pi} + \frac{1}{2^n} e^{i \frac{1}{3} \pi} + \sum_{k=n+1}^{\infty} \frac{1}{2^k} e^{i \frac{2^{k-n}}{3} \pi} \\ &= \sum_{k=1}^{n-1} \frac{1}{2^k} e^{i \frac{2^k}{2^n 3} \pi} + \frac{1}{2^n} \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) + \frac{1}{2^n} \sum_{k=1}^{\infty} \frac{1}{2^k} e^{i \frac{2^k}{3} \pi}. \end{aligned}$$

Notice that $e^{i \frac{2^k}{3} \pi} = e^{i(-1)^{k+1} \frac{2}{3} \pi} = -\frac{1}{2} + i(-1)^{k+1} \frac{\sqrt{3}}{2}$. Thus, $\sum_{k=1}^{\infty} \frac{1}{2^k} e^{i \frac{2^k}{3} \pi} = -\frac{1}{2} + i \frac{\sqrt{3}}{6}$. After these computations $c_{2^n 3}$ becomes

$$\sum_{k=1}^{n-1} \frac{1}{2^k} e^{i \frac{2^k}{2^n 3} \pi} + i \frac{2\sqrt{3}}{2^n 3}.$$

Now using the inequality $\cos(\alpha) \leq 1 - \frac{\alpha^2}{2} + \frac{\alpha^4}{24}$ we obtain the following estimate

$$\begin{aligned} |\Re c_{2^n 3}| &\leq \sum_{k=1}^{n-1} \frac{1}{2^k} \left(1 - \frac{\pi^2 4^k}{18 \cdot 4^n} + \frac{\pi^4}{81 \cdot 24} \frac{16^k}{16^n} \right) \\ &= 1 - \frac{1}{2^{n-1}} - \frac{\pi^2}{18} \frac{2^n - 2}{4^n} + \frac{\pi^4}{81 \cdot 24} \frac{8^n - 8}{16^n 7} \\ &= 1 - \frac{1}{2^n 3} \left(6 + \frac{\pi^2}{6} - \frac{\pi^4}{27 \cdot 24 \cdot 7} \right) + \frac{a}{(2^n 3)^2} + \frac{b}{(2^n 3)^4}, \end{aligned}$$

for some constants a and b . Similarly, using $\sin(\alpha) \leq \alpha$, we have

$$|\Im c_{2^n 3}| \leq \sum_{k=1}^{n-1} \frac{1}{2^k} \frac{2^k}{2^n 3} \pi + \frac{2\sqrt{3}}{2^n 3} \leq \frac{(n+1)\pi}{2^n 3}.$$

Thus,

$$\begin{aligned} |c_{2^n 3}|^{2^n 3} &= (|\Re c_{2^n 3}|^2 + |\Im c_{2^n 3}|^2)^{\frac{2^n 3}{2}} \\ &\leq \left(1 - \frac{2}{2^n 3} \left(6 + \frac{\pi^2}{6} - \frac{\pi^4}{27 \cdot 24 \cdot 7} \right) + \left(\frac{2}{2^n 3} \right)^2 (n+1)^2 a_1 \right. \\ &\quad \left. + \left(\frac{2}{2^n 3} \right)^2 a_2 + \dots + \left(\frac{2}{2^n 3} \right)^8 a_8 \right)^{\frac{2^n 3}{2}}. \end{aligned}$$

Passing to the upper limit as $n \rightarrow \infty$, we finally obtain (4).

2.2. L_p -case. Our second example is on the Hilbert space $L_2[0, 2\pi]$, but now for a positive contractive semigroup and positive contractive projection.

We take $e^{tA} f(x) = f(x + 2\pi t)$, regarding f as a 2π -periodic function. Now let P be the orthogonal projection onto the space spanned by the

positive norm-one function $g(x) = \frac{1}{\sqrt{34\pi}} \left[4 + \sum_{k=0}^{\infty} \frac{1}{\sqrt{2^k}} \cos 2^k x \right]$. Notice that like in the previous example our projection is one-dimensional (see Remark 1 below). Simple substitution shows that (1) evaluated at g for $t = 1$ exists if and only if the numerical limit

$$\lim_{n \rightarrow \infty} \left[\int_0^{2\pi} g(x) g\left(x + \frac{1}{n}\right) dx \right]^n$$

exists. Denoting

$$c_n = \int_0^{2\pi} g(x) g\left(x + \frac{1}{n}\right) dx$$

and using the orthogonality of cosines, we get

$$c_n = \frac{16}{17} + \frac{1}{17} \sum_{k=1}^{\infty} \frac{1}{2^k} \cos \frac{2^k}{n} \pi$$

Now following the same calculations as for the first example, we obtain inequalities (3) and (4) with powers doubled on the right hand sides.

This disproves the second conjecture.

Remark 1. As we have already noticed the projections in our examples are one-dimensional. It would be interesting to know what property of a semigroup on a Hilbert space is responsible for the existence of (1) for all one-dimensional, or more specifically, one-dimensional orthogonal projections.

The authors are grateful to Wolfgang Arendt, András Bátkai and Bálint Farkas for helpful conversations.

REFERENCES

- [1] W. Arendt and M. Ulm, *Trotter's product formula for projections*, (1997), http://www.mathematik.uni-ulm.de/ees/pr_archi.html.
E-mail address: matomate@cs.elte.hu

ELTE TTK, DEPARTMENT OF APPLIED ANALYSIS, 1053 BUDAPEST, KECSKEMÉTI U. 10-12, HUNGARY.

E-mail address: shvidkoy@math.missouri.edu

MATHEMATICS DEPARTMENT, UNIVERSITY OF MISSOURI-COLUMBIA, COLUMBIA, MO 65211, USA.